

AD-A110 454

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/S 20/4

NEW DESCRIPTIONS OF DISPERSION IN FLOW THROUGH TUBES: CONVOLUTION--ETC(U)

NOV 81 J C WANG, W E STEWART

DAAG29-80-C-0041

UNCLASSIFIED


MRC-TSR-2297

NL

1 1 1  
20 10 20 10

1

END  
DATE  
FILMED  
3 82  
DTIC



**LEVEL**

②

AD A110454

MRC Technical Summary Report #2297

NEW DESCRIPTIONS OF DISPERSION IN  
FLOW THROUGH TUBES:  
CONVOLUTION AND COLLOCATION METHODS

James C. Wang and Warren E. Stewart

Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706

November 1981

Received August 5, 1981)

DTIC  
COLLECTED  
FEB 4 1982  
D  
H

Approved for public release  
Distribution unlimited

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

and

National Science Foundation  
Washington, DC 20550

82 02 03 085

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

NEW DESCRIPTIONS OF DISPERSION IN FLOW THROUGH TUBES:  
CONVOLUTION AND COLLOCATION METHODS

James C. Wang and Warren E. Stewart

Technical Summary Report #2297

November 1981

ABSTRACT

The convective dispersion of a solute in steady flow through a tube is analyzed, and the concentration profile for any Peclet number is obtained as a convolution of the profile for infinite Peclet number. Close approximations are obtained for the concentration profile and its axial moments, by use of orthogonal collocation in the radial direction. The moments thus obtained converge rapidly, and the concentration profile less rapidly, toward exactness as the number of collocation points is increased. A two-point radial grid gives results of practical accuracy; analytical solutions are obtained at this level of approximation.

AMS (MOS) Subject Classifications: 35A35, 41A10, 41A63, 44A30, 65N35, 65N40,  
76R05

Key Words: Dispersion, Diffusion, Polynomial approximation, Orthogonal  
collocation, Method of lines, Laplace transform, Fourier  
transform, Convolution, Superposition, Moment-generating function

Work Unit Number 2 - Physical Mathematics

---

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041 and  
the National Science Foundation under Grant Numbers ENG76-24368 and CPE79-  
13162.


## SIGNIFICANCE AND EXPLANATION

Convective dispersion plays an important role in many processes of chemical reaction and separation. Such systems are commonly analyzed by use of a radially-averaged diffusion equation. This approach has been popularized by the simple results of Taylor (1953) and Aris (1956) for the cross-sectional average concentration, and perhaps also by the belief that the radial variations of concentration are too small to require detailed consideration. Subsequent investigators (e.g. Gill and Sankarasubramanian (1970), Chatwin (1977), De Gance and Johns (1978)) have extended this approach to shorter times and to other boundary conditions.

The foregoing approach is not easy to apply to reaction or separation processes if the fluid properties vary, nor if the kinetics or equilibria are non-linear. Therefore, in this paper we consider an alternate method: we solve the full diffusion equation by orthogonal collocation in the radial direction. Non-reactive systems are emphasized here; reactive ones will be studied more fully in the sequel to this p.

Accession For DTIC DTIC DTIC DTIC	<input checked="" type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	By Distribution/ Availability Codes and/or Social
<div style="font-size: 48px; font-family: cursive;">A</div>		

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the authors of this report.



# NEW DESCRIPTIONS OF DISPERSION IN FLOW THROUGH TUBES: CONVOLUTION AND COLLOCATION METHODS

James C. Wang and Warren E. Stewart

## INTRODUCTION

The collocation procedure described here is a variant of that given by Villadsen and Stewart (1967); see also Finlayson (1972) and Villadsen and Michelsen (1978). For axially symmetric states in a tubular reactor, we approximate the profile of each unknown state variable  $Y_m$  in the form

$$\tilde{Y}_m = \sum_{i=0}^n a_{im}(\tau, Z) \xi^{2i} \quad 0 \leq \xi \leq 1 \quad (1)$$

This expansion can also be written as a Lagrange polynomial

$$\tilde{Y}_m = \sum_{k=1}^{n+1} w_k(\xi) \tilde{Y}_m(\tau, Z, \xi_k) \quad 0 \leq \xi \leq 1 \quad (2)$$

with radial basis functions

$$w_k(\xi) = \prod_{j \neq k} (\xi^2 - \xi_j^2) / \prod_{j \neq k} (\xi_k^2 - \xi_j^2) \quad (3)$$

Thus, each approximate profile  $\tilde{Y}_m$  is represented by Lagrange interpolation in terms of its values  $\tilde{Y}_m(\tau, Z, \xi_k)$  at a set of chosen radial nodes  $\xi_k$ . We choose  $\xi_1^2, \dots, \xi_n^2$  as the zeros of the shifted Legendre polynomial  $P_n(x)$  (Abramowitz and Stegun, 1972) and choose  $\xi_{n+1}$  as the wall location,  $\xi = 1$ .

Any needed derivatives or integrals of  $\tilde{Y}_m$  can now be represented as linear combinations of the nodal values  $\tilde{Y}_m(\xi_k)$ ; this is known as the method of ordinates (Villadsen and Stewart, 1967). For example, the dimensionless radial derivatives of interest at the radial node  $\xi_i$  are:

$$\begin{aligned} \left. \frac{d\tilde{Y}_m}{d\xi} \right|_{\tau, Z, \xi_i} &= \sum_{k=1}^{n+1} \left( \frac{d}{d\xi} w_k(\xi) \right) \Big|_{\xi_i} \tilde{Y}_m(\tau, Z, \xi_k) \\ &= \sum_{k=1}^{n+1} A_{ik} \tilde{Y}_{mk}(\tau, Z) \end{aligned} \quad (4)$$

$$\begin{aligned}
\frac{1}{\xi} \frac{d}{d\xi} \left( \xi \frac{d\tilde{Y}_m}{d\xi} \right) \Big|_{\tau, Z, \xi_i} &= \sum_{k=1}^{n+1} \left[ \frac{1}{\xi} \frac{d}{d\xi} \left( \xi \frac{dW_k}{d\xi} \right) \Big|_{\xi_i} \right] \tilde{Y}_m(\tau, Z, \xi_k) \\
&= \sum_{k=1}^{n+1} B_{ik} \tilde{Y}_{mk}(\tau, Z) \quad .
\end{aligned}
\tag{5}$$

In this way, we reduce all radial derivatives to summations; the same can be done for differences and integrals with respect to  $\xi$ . For the above choice of nodes, weighted integrals of  $\tilde{Y}_m$  over the cross-section can be obtained with high accuracy by use of Gauss' quadrature formula.

The boundary conditions at the wall may be linear or non-linear. If they are linear, then  $\tilde{Y}_{m,n+1}$  can be expressed linearly in terms of the interior values  $\tilde{Y}_{mk}$  by use of the boundary condition and Eq. (4). This permits elimination of  $\tilde{Y}_{m,n+1}$  from Eq. (5); the resulting new weight coefficients will be denoted by  $B'_{ik}$ . In non-linear cases, on the other hand, one retains the wall values as working variables and applies the boundary conditions numerically during the solution process.

#### CONVOLUTION RELATION

Consider the developed isothermal flow of a binary Newtonian or non-Newtonian fluid in an infinitely long circular tube. The continuity equation for either species may be written in the dimensionless form

$$\frac{\partial \Omega}{\partial \tau} + v(\xi) \frac{\partial \Omega}{\partial Z} = \frac{1}{\xi} \frac{\partial}{\partial \xi} \left( \xi \frac{\partial \Omega}{\partial \xi} \right) + \frac{1}{Pe^2} \frac{\partial^2 \Omega}{\partial Z^2} - K'' \Omega \tag{6}$$

with the initial and boundary conditions

$$\text{At } \tau = 0 : \Omega(\tau, Z, \xi) = G(Z, \xi) \tag{7}$$

$$\text{At } \xi = 1 : \frac{\partial}{\partial \xi} \Omega(\tau, Z, \xi) + K'' \Omega(\tau, Z, \xi) = 0 \text{ for } \tau > 0 \tag{8}$$

$$\text{At } \xi = 0 : \frac{\partial}{\partial \xi} \Omega(\tau, Z, \xi) = 0 \text{ for } \tau > 0 . \quad (9)$$

Here  $K'''$  and  $K''$  are the first and second dimensionless numbers of Damköhler (1936) for homogeneous and heterogeneous reactions. The Laplace-Fourier transform of Eqs. (6) - (9) from  $(\tau, Z, \xi)$  into  $(s, p, \xi)$  is

$$\left(s + K''' - \frac{p^2}{Pe^2}\right)\bar{\Omega} + pV(\xi)\bar{\Omega} = \frac{1}{\xi} \frac{\partial}{\partial \xi} \left(\xi \frac{\partial \bar{\Omega}}{\partial \xi}\right) + \bar{G}(p, \xi) \quad (10)$$

$$\text{At } \xi = 1 : \frac{\partial}{\partial \xi} \bar{\Omega}(s, p, \xi) + K''\bar{\Omega}(s, p, \xi) = 0 \quad (11)$$

$$\text{At } \xi = 0 : \frac{\partial}{\partial \xi} \bar{\Omega}(s, p, \xi) = 0 . \quad (12)$$

Let  $\Omega_\infty$  be the solution of Eqs. (6) - (9) with  $Pe = \infty$ . Its transform satisfies eqs. (10) - (12) with the  $p^2$  term suppressed. Use of the s-shift theorem and Fourier convolution (Campbell and Foster 1967) then gives:

$$\bar{\Omega}(\tau, p, \xi) = \exp\left(\frac{p^2 \tau}{Pe^2}\right) \bar{\Omega}_\infty(\tau, p, \xi) \quad (13)$$

$$\Omega(\tau, Z, \xi) = \int_{-\infty}^{\infty} \frac{Pe}{2\sqrt{\pi\tau}} \exp\left[-\frac{Pe^2}{4\pi\tau}(Z-X)^2\right] \Omega_\infty(\tau, X, \xi) dX . \quad (14)$$

Thus, the complete solution of Eqs. (6) - (9) is obtained by Gaussian smoothing of the infinite-Pe solution.

#### TWO-POINT COLLOCATION; NON-REACTIVE CASE

Let  $C_\infty$  be the solution of Eqs. (6) - (9) with  $K'' = K''' = 0$ ,  $Pe = \infty$ , and the following initial mass fraction profile:

$$G(Z, \xi) = \delta(Z)\phi(\xi) . \quad (15)$$

Collocation with  $n = 2$ , and elimination of the wall concentration through the boundary condition (8), yields an initial-value problem for the nodal



functions  $\tilde{C}_\infty(\tau, Z, \xi_1) \equiv \tilde{C}_{1\infty}(\tau, Z)$  and  $\tilde{C}_\infty(\tau, Z, \xi_2) \equiv \tilde{C}_{2\infty}(\tau, Z)$ . The boundary condition (9), and the symmetry of the true solution, are satisfied through the use of only even powers of  $\xi$  in the collocation basis functions [see Eq. (3)]. In place of Eq. (10) we then obtain

$$s\tilde{C}_{i\infty} + pV_i\tilde{C}_{i\infty} = \sum_{k=1}^2 B'_{ik}\tilde{C}_{k\infty} + Q_i \quad i = 1, 2 \quad (16)$$

in which  $V_i$  and  $Q_i$  are the nodal values of  $V(\xi)$  and  $Q(\xi)$ . To complete the solution, we evaluate the coefficients  $B'_{ik}$  from Eqs. (5) and (8), with the Gaussian nodes

$$\left. \begin{aligned} \xi_1^2 &= \frac{1}{2} - \frac{1}{\sqrt{12}} \\ \xi_2^2 &= \frac{1}{2} + \frac{1}{\sqrt{12}} \end{aligned} \right\} \quad (17)$$

and invert the transforms. The solution differs from zero only within the region  $|\tilde{Z}| < |T|$ , and is given there by

$$C_{1\infty} = \exp(-8\tau) \left\{ \frac{Q_1 \beta |T+\tilde{Z}| I_1(x)}{2x} + \frac{8Q_2 I_0(x)}{|V_1 - V_2|} + Q_1 \delta(\tilde{Z}-T) \right\} \quad (18.1)$$

$$C_{2\infty} = \exp(-8\tau) \left\{ \frac{Q_2 \beta |T-\tilde{Z}| I_1(x)}{2x} + \frac{8Q_1 I_0(x)}{|V_1 - V_2|} + Q_2 \delta(\tilde{Z}+T) \right\} \quad (18.2)$$

in which

$$\tilde{Z} = Z - (V_1 + V_2)\tau/2 \quad (19)$$

$$T = (V_1 - V_2)\tau/2 \quad (20)$$

$$\beta = 256(V_1 - V_2)^{-2} \quad (21)$$

$$x = \sqrt{\beta(T^2 - \tilde{Z}^2)} \quad (22)$$

Interpolating the solution according to Eq. (2), and integrating over the tube cross-section, we find the mean value to be  $(C_{1\infty} + C_{2\infty})/2$ . Gaussian quadrature of the integral gives the same result.

#### CENTRAL MOMENTS OF THE TWO-POINT SOLUTION

Let  $C$  be the solution of Eqs. (6) - (9) and (15) with  $K'' = K''' = 0$ . The mean mass fraction in a cross-section is defined, for these axisymmetric problems, by

$$\langle C \rangle = 2 \int_0^1 C \xi d\xi \quad (23)$$

The central axial moments of  $\langle C \rangle$  are

$$\mu_i = \int_{-\infty}^{\infty} (Z - \langle V \rangle \tau)^i \langle C \rangle dZ \quad (24)$$

and have the moment-generating function

$$\mu_i = \left( -\frac{d}{dp} \right)^i [\exp(\langle V \rangle \tau p) \langle \bar{C}(\tau, p, \xi) \rangle] \Big|_{p=0} \quad (25)$$

in which  $\bar{C}(\tau, p, \xi)$  is the Fourier transform of  $C(\tau, Z, \xi)$ .

For the collocation solution with  $n = 2$ , we thus obtain

$$\tilde{\mu}_0 = \frac{1}{2} (Q_1 + Q_2) \quad (26)$$

$$\tilde{\mu}_1 = (Q_1 - Q_2)(V_1 - V_2) \left( \frac{1 - \exp(-16\tau)}{64} \right) \quad (27)$$

$$\tilde{\mu}_2 = \tilde{\mu}_0 \left\{ \frac{2\tau}{Pe^2} + \frac{(V_1 - V_2)^2}{8} \left( \frac{\tau}{4} - \frac{1 - \exp(-16\tau)}{64} \right) \right\} \quad (28)$$

For a radially uniform pulse of unit strength,  $Q_i = 1$ ; and for developed laminar flow of a Newtonian fluid,  $V_i = 1 - \xi_i^2$ . Eqs. (26) - (28) then give

$$\tilde{\mu}_0 = 1 \quad (29)$$

$$\tilde{\mu}_1 = 0 \quad (30)$$

$$\tilde{\mu}_2 = \left( \frac{1}{Pe^2} + \frac{1}{192} \right) 2\tau - \frac{1 - \exp(-16\tau)}{1536} \quad (31)$$

The exact solutions derived by Aris (1956), Gill and Sankarasubramanian (1970), and Chatwin (1977) for a radially uniform pulse confirm Eqs. (29) and (30). Their expressions for  $\mu_2$  can be reduced (after a sign correction in Aris' solution) to the common form

$$\mu_2 = \left(-\frac{1}{Pe} + \frac{1}{192}\right) 2\tau - \frac{1}{1440} + 32 \sum_{j=1}^{\infty} \lambda_j^{-8} \exp(-\lambda_j^2 \tau) \quad (32)$$

in which  $\lambda_j$  is the  $j$ th zero of the Bessel function  $J_1(x)$ . The terms proportional to  $\tau$  constitute the prediction of the standard Fickian dispersion model, with the long-time asymptotic dispersion coefficient found by Taylor (1953) and completed by Aris (1956). For brevity, the latter model is called the Taylor-Aris model.

Before testing these results numerically, we give collocation solutions of higher order for the axial moments of the function  $C(\tau, Z, \xi)$ .

#### COLLOCATION SOLUTIONS OF HIGHER ORDER

The moments of the mass fraction profile about the origin are defined by

$$m_1(\tau, \xi) = \int_{-\infty}^{\infty} Z^1 \Omega(\tau, Z, \xi) dZ \quad (33)$$

The generating function for these moments is

$$m_1(\tau, \xi) = \left(-\frac{d}{dp}\right)^1 [\bar{\Omega}(\tau, p, \xi)] \big|_{p=0} \quad (34)$$

The moments at finite  $Pe$  can be obtained from those at infinite  $Pe$  by use of Eqs. (13) and (34).

A collocation solution for  $m_1(\tau, \xi)$  at  $Pe = \infty$  is obtained as follows.

Let  $\underline{m}_1$  be the vector of mesh-point values of  $m_1(\tau, \xi)$ :

$$\underline{m}_1 = [m_1(\tau, \xi_1), \dots, m_1(\tau, \xi_n)]^T \quad (35)$$

We apply the Fourier transformation and  $n$ -point radial orthogonal collocation to Eqs. (6) - (9) with  $K'' = K''' = 0$  and  $Pe = \infty$ . Then by use of Eq. (34)

we obtain

$$\frac{\partial}{\partial \tau} m_i = B' m_i + i \sum m_{i-1} \quad (36)$$

Eqs. (13), (34), and (36) can then be used to obtain the moments at finite values of  $\tau$ . Finally, the cross-sectional average  $\langle m_i \rangle$  can be obtained by Gaussian quadrature of  $m_i$ .

Tables 1, 2, and 3 show the collocation solutions and the exact solutions for  $\mu_0$ ,  $\mu_1$ , and  $\mu_2$  for two initial solution distributions. The collocation solutions converge rapidly toward the exact results for all values of  $\tau$ , and for both initial distributions. From Table 1 we see that the collocation solution for  $n = 2$  is particularly accurate in the case of a uniform initial radial distribution.

#### TWO-DIMENSIONAL INITIAL DISTRIBUTIONS

Any axially symmetric initial distribution can be expressed in the form

$$G(Z, \xi) = \sum_j \sum_k b_{jk} \varphi^j(Z) \varrho^k(\xi) \quad (37)$$

with suitable basis functions  $\varphi^j(Z)$  and  $\varrho^k(\xi)$ . Let  $C^k(\tau, Z, \xi)$  be the solution of Eqs. (6) - (9) with constant  $K''$  and  $K'''$ , and with the initial distribution  $C^k(0, Z, \xi) = \delta(Z) \varrho^k(\xi)$ . Then the solution corresponding to the initial condition (37) is obtainable by superposition,

$$\Omega(\tau, Z, \xi) = \sum_j \sum_k b_{jk} \int_{-\infty}^{\infty} \varphi^j(x) C^k(\tau, Z-x, \xi) dx \quad (38)$$

and has the Fourier transform

$$\bar{\Omega}(\tau, p, \xi) = \sum_j \sum_k b_{jk} \bar{\varphi}^j(p) \bar{C}^k(\tau, p, \xi) \quad (39)$$

Let  $\phi_i^j$  and  $M_i^k$  be the  $i$ th moments of the functions  $\varphi^j(Z)$  and  $C^k(\tau, Z, \xi)$  respectively with respect to  $Z$ . Then from Eqs. (34) and (39) we get the moments  $m_i$  in the form

Table 1 Second moment  $1000(\mu_2 - \frac{2\tau}{Pe^2})$  for initial condition  $\phi(\xi) = 1$

$\tau$	Collocation			Exact *	Taylor-Aris
	n=2	n=3	n=4		
.01	.00791	.00782	.00780	.00779	.1042
.05	.1623	.1575	.1574	.1574	.5208
.10	.5221	.5059	.5059	.5059	1.042
.15	.9705	.9441	.9442	.9442	1.562
.20	1.459	1.425	1.425	1.425	2.083
.25	1.965	1.927	1.927	1.927	2.604
.40	3.517	3.474	3.474	3.474	4.167
1.0	9.766	9.722	9.722	9.722	10.42

\* Exact values are calculated from Eq. (32). The first two moments agree for all three methods.

Table 2 First moment  $100 \mu_1$  for initial condition  $\phi(\xi) = 2\xi^2$

$\tau$	Collocation			Exact*
	n=2	n=3	n=4	
.01	-.1540	-.1516	-.1511	-.1511
.05	-.5736	-.5540	-.5541	-.5541
.10	-.8314	-.8083	-.8086	-.8086
.15	-.9472	-.9298	-.9299	-.9299
.20	-.9992	-.9880	-.9880	-.9880
.25	-1.0226	-1.0160	-1.0159	-1.0159
.40	-1.0399	-1.0388	-1.0388	-1.0388
1.0	-1.0417	-1.0417	-1.0417	-1.0417

\*Exact values are calculated from Chatwin (1977). The first moment is independent of  $Pe$ , in view of Eqs. (13) and (34).

Table 3 Second moment  $1000(\mu_2 - \frac{2\tau}{Pe^2})$  for initial condition  $\phi(\xi) = 2\xi^2$

$\tau$	Collocation			Exact*
	n=2	n=3	n=4	
.01	.00791	.00747	.00738	.00737
.05	.1623	.1390	.1389	.1389
.10	.5221	.4434	.4447	.4447
.15	.9705	.8415	.8431	.8431
.20	1.459	1.294	1.296	1.296
.25	1.965	1.778	1.779	1.779
.40	3.517	3.305	3.305	3.305
1.0	9.766	9.549	9.549	9.549

\*Exact values are calculated from Chatwin (1977).

$$m_i(\tau, \xi) = \sum_j \sum_k b_{jk} \sum_{r=0}^i \binom{i}{r} \phi_{i-r}^j m_r^k(\tau, \xi) \quad (40)$$

for any initial distribution of the form in Eq. (37). This result shows the influence of the initial conditions on the moments of the mass fraction profile at any later time.

#### PROFILES FOR A PULSE OF FINITE LENGTH

As a further test of the collocation solutions, we calculate the mass fraction profiles for an initial pulse of maximum amplitude unity and of finite length  $L$  equal to 0.1 unit of  $Z$ . The coordinate

$$U = (2Z - \tau)/(L + \tau) \quad (41)$$

is introduced here to facilitate finite element calculations of the solution. The following initial condition is used,

$$G(Z, \xi) = \varphi(U) \quad (42)$$

in which  $\varphi(U)$  is a  $C^1$  cubic spline:

$$\left. \begin{aligned} \varphi(U) &= 1 && \text{for } |U| < 0.975 \\ \varphi(U) &= 1 - 3 \left( \frac{|U| - 0.975}{0.025} \right)^2 + 2 \left( \frac{|U| - 0.975}{0.025} \right)^3 && \text{for } 0.975 < |U| < 1 \\ \varphi(U) &= 0 && \text{for } |U| > 1 \end{aligned} \right\} \quad (43)$$

With this initial condition, the solution  $C_\infty$  (valid for  $Pe = \infty$ ) is permanently zero outside the region  $-1 < U < 1$ .

Figures 1, 2, and 3 show the cross-sectional average mass fraction  $\langle w \rangle$  as a function of  $\tau$  and  $z$  at  $Pe = 15$ , 40, and infinity. Three methods of solution are shown: orthogonal collocation on finite elements of  $U$ , two-point radial collocation, and the Taylor-Aris model. The radial collocation was done in all cases as described above; the finite elements of  $U$  were  $C^1$  cubic splines with length  $\Delta U = 0.025$ . Each method was first applied with



$Pe = \infty$ , and Eq. (14) was then used to get results at finite Peclet numbers.

The two-point collocation profiles (dashed curves) model the true solutions very well, except for some lack of smoothness at short times with  $Pe = \infty$ .

The Taylor-Aris model exaggerates the width of the tails of the pulse in every case.




Figure 1. Radially averaged profiles at  $Pe = 15$ , for the initial condition of Eqs. (42)-(43).

— Reference solution by finite elements  
 --- Collocation solution with  $n = 2$   
 .... Solution by Taylor-Aris model




Figure 2. Radially averaged profiles at  $Pe = 40$ , for the initial condition of Eqs. (42)-(43)

— Reference solution by finite elements  
 --- Collocation solution with  $n = 2$   
 .... Solution by Taylor-Aris model




Figure 3. Radially averaged profiles at  $Pe = \infty$ , for the initial condition of Eqs. (42)-(43)

— Reference solution by finite elements  
 --- Collocation solution with  $n = 2$   
 .... Solution by Taylor-Aris model

# NOTATION

$A_{ik}$  coefficients for gradient operator in Eq. (4)

$a_{im}, b_{jk}$  coefficients of basis functions.

$B_{ik}$  coefficients for Laplacian operator in Eq. (5)

$B'_{ik}$  coefficients for Laplacian operator with boundary point eliminated.

$B'$  matrix of order  $n$  with elements  $B'_{ik}$

$C$  solution of Eqs. (6) - (9) with constant  $K''$  and  $K'''$  and with initial condition  $\delta(z)\phi(\xi)$

$C^k$  special form of  $C$  for initial condition  $\delta(z)\phi^k(\xi)$

$C_\infty$  limit of  $C$  as  $Pe \rightarrow \infty$

$G(\xi, z)$  initial distribution, Eq. (7)

$I_n(x)$  modified Bessel function of the first kind

$K''', K''$  first and second Damköhler numbers

$m_i$   $i$ th axial moment of  $\Omega$  defined in Eq. (33)

$\underline{m}_i$  vector of nodal values of  $m_i$ , Eq. (35)

$n$  number of collocation points interior to  $\xi = 1$

$p$  Fourier transform variable

$Pe = RV_{\max}/D_{AB}$ , Peclet number

$Q(\xi)$  radial function in initial condition

$Q_i$  value of  $Q(\xi)$  at  $i$ th collocation point

$Q^k(\xi)$   $k$ th radial basis function in initial condition

$s$  Laplace transform variable

$U$  transformed  $Z$  coordinate in Eq. (41)

$V(\xi)$  velocity profile, relative to centerline value

$V_i$  value of  $V(\xi)$  at  $i$ th collocation point

$V$  diagonal matrix with elements  $V_{ii} = V_i$

$Y_m$  mth state variable (e.g., temperature or species mass fraction)

$Z = z D_{AB} / (V_{\max} R^2)$ , dimensionless axial coordinate

$\langle \rangle$  average over flow cross-section

$\binom{i}{r}$  binomial coefficient

#### Greek Letters

$\mu_i$  ith central moment of  $C$  defined in Eq. (24)

$\xi = r/R$ , dimensionless radial coordinate

$\tau = t D_{AB} / R^2$ , dimensionless time

$\varphi(U)$  axial function in initial condition of Eq. (43)

$\varphi^j(z)$  jth axial basis function in initial condition of Eq. (37)

$\omega$  solute mass fraction

$\Omega$  solution for  $\omega$  with general initial condition  $G(z, \xi)$

#### LITERATURE CITED

- Abramowitz, M. and I. A. Stegun, ed., "Handbook of Mathematical Functions", Dover (1972).
- Aris, R., "On the Dispersion of a Solute in a Fluid Flowing Through a Tube", Proc. Roy. Soc. A, 235, 67 (1956).
- Campbell, G. A. and R. M. Foster, "Fourier Integrals for Practical Applications", D. Van Nostrand, Princeton, New Jersey (1967).
- Chatwin, P. C., "The Initial Development of Longitudinal Dispersion in Straight Tubes", J. Fluid Mech., 80, 33 (1977).
- Danköbler, G., "Einfluss der Strömung, Diffusion und Wärmeüberganges auf die Leistung von Reaktionsöfen. I. Allgemeine Gesichtspunkte für die Übertragung eines chemischen Prozesses aus dem Kleinen ins Grosse", Z. Electrochem. 42, 846 (1936).
- De Gance, A. E. and L. E. Johns, "On the Dispersion Coefficients for Poiseuille Flow in a Circular Cylinder", Appl. Sci. Res., 34, 227 (1978).
- Finlayson, B. A., "The Method of Weighted Residuals and Variational Principles", Academic Press, New York (1972).
- Gill, W. N. and R. Sankarasubramanian, "Exact Analysis of Unsteady Convective Diffusion", Proc. Roy. Soc. A, 316, 341 (1970).
- Taylor, Sir Geoffrey, "Dispersion of Soluble Matter in Solvent Flowing Slowly Through a Tube", Proc. Roy. Soc. A, 219, 186 (1953).
- Villadsen, J. V. and M. L. Michelsen, "Solutions of Differential Equation Models by Polynomial Approximation", Prentice-Hall, Englewood Cliffs, New Jersey (1978).
- Villadsen, J. V. and W. E. Stewart, "Solution of Boundary-Value Problems by Orthogonal Collocation", Chem. Eng. Sci., 22, 1483 (1967), 23, 1515 (1968).

JCW/WES/jvs

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER #2297	2. GOVT ACCESSION NO. AD-A110454	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  New Descriptions of Dispersion in Flow Through Tubes: Convolution and Collocation Methods		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s)  James C. Wang and Warren E. Stewart		8. CONTRACT OR GRANT NUMBER(s) ENG76-24368; CPE79-13162 DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 2 - Physical Mathematics
11. CONTROLLING OFFICE NAME AND ADDRESS  (see Item 18 below)		12. REPORT DATE November 1981
		13. NUMBER OF PAGES 18
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)  UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES U. S. Army Research Office and National Science Foundation P. O. Box 12211 Washington, DC 20550 Research Triangle Park North Carolina 27709		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Dispersion, Diffusion, Polynomial approximation, Orthogonal collocation, Method of lines, Laplace transform, Fourier transform, Convolution, Superposition, Moment-generating function		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The convective dispersion of a solute in steady flow through a tube is analyzed, and the concentration profile for any Peclet number is obtained as a convolution of the profile for infinite Peclet number. Close approximations are obtained for the concentration profile and its axial moments, by use of orthogonal collocation in the radial direction. The moments thus obtained converge rapidly, and the concentration profile less rapidly, toward exactness as the number of collocation points is increased. A two-point radial grid gives results of practical accuracy; analytical solutions are obtained at this level of approximation.		



**DAT  
FILM**